

PARRY SOUND AREA LAKE SUMMARIES

Brought to you compliments of: Gord Pollock, Broker Century 21 Granite Properties Ltd Brokerage 33 James Street, Parry Sound, Ontario P2A 1T6 Business: 705-746-2158, Fax: 705-746-4746 Residence: 705-746-4816, Cell: 705-774-0955

Website: www.gordpollock.com Email: gord@gordpollock.com

Squaw Lake

Created: November 03 Revised:

Location:

MNR District:	Parry Sound
Geographic Township:	Blair
Municipal Township:	Unincorporated
Watershed:	Pickerel River
Angling Division:	15

Basin and Terrain Characteristics:

Lake Survey Year:	1977
Surface Area:	132.9 hectares
Maximum Depth:	20.1 meters
Mean Depth:	4.1 meters
Perimeter:	14.7 km
Island shoreline:	0.8 km
Littoral Zone:	
Thermal Regime:	Cool
Shoreline Development:	
Access Points:	Road (Public)
Water Level:	Not Regulated
Crown Land:	100% shoreline

Water Quality:

(Parameters pertain to fisheries habitat only. For information on potability of water or contaminates, contact Min. of Health and Min. of Environment.)

Secchi reading: 2.5 meters Colour: Yellow/Brown Dissolved Oxygen:Alkalinity:13.7 mg/l - Level 3 Moderately Sensitive (MoE, 1989)pH:7.0Total Phosphorus:7.6

"Guide to eating fish": No Information

Fisheries:

- Game Fish Species: Walleye (1990), Smallmouth Bass (1990), Largemouth Bass (1990), Muskellunge (1990), Northern Pike, Yellow Perch
- Other species present: Rock Bass (1990), Pumpkinseed (1990), Northern Redhorse Sucker (1990), Channel Catfish (1990), Bowfin (1990), Carp (1990), Silver Lamprey, Lake Whitefish, Emerald Shiner, Mimic Shiner, White Sucker, Bluntnose Minnow, Brown Bullhead, Burbot, Johnny Darter, Logperch

Exotic Species:

Stocking Record:

Stress Type:

Use Type: Recreation, Recreational Fishing, Canoe Route

Summary of Fisheries Studies / Reports:

Emerson, R. & Thurston, L. 1992 1990 Trap Net Survey on Squaw lake (Blair Township)

- trapnets utilized were 8, 6 and 4 foot for a total of 15 sets
- The C.U.E. for walleye 0.4; smallmouth bass 0.8; muskellunge 0.4 indicating the walleye and smallmouth bass populations are low and the muskellunge population can be considered fair.
- The C.U.E. for largemouth bass (2.3) with the 4 foot nets indicates the population is fair.
- The bulk of the catch (67%) was made up of the two coarse fish catfish and redhorse suckers.

MNR 1977 Raw Data on File: Creel Survey 1974, 76 & 77

- 16 anglers interviewed with 48.75 angler hours having caught 19 northern pike and 9 bass
- Species sought included northern pike, bass, walleye and muskellunge

Management Prescription: